



## New polymorphic microsatellite markers for the endangered fern *Ceratopteris thalictroides* (Parkeriaceae)

Won-Bum CHO, Eun-Kyeong HAN, Myounghai KWAK<sup>1</sup> and Jung-Hyun LEE\*

*Department of Biology Education, Chonnam National University, Gwangju 61186, Korea*

<sup>1</sup>*Plant Resources Division, National Institute of Biological Resources, Incheon 22689, Korea*

(Received 31 May 2018; Revised 14 June 2018; Accepted 18 June 2018)

**ABSTRACT:** *Ceratopteris thalictroides* is a semi-aquatic fern with a circumtropical distribution. Because this species is designated internationally on the IUCN Red List as requiring at least some concern, Korean populations are of great concern for the species' long-term survival, as they are at the northern limit of the species distribution. To establish an effective conservation strategy for those populations at the genetic level, we used the Mi-Seq platform to develop three sets of 25 polymorphic microsatellite markers for *C. thalictroides*, which is endangered in Korea. In populations sampled from Busan and Gochang, the number of alleles ranged from 2 to 13 (average of 5.64), and plants presented an expected heterozygosity of 0.000 to 0.860. These markers will be useful for evaluating the genetic status and conserving Korean populations of *C. thalictroides* more effectively.

**Keywords:** *Ceratopteris thalictroides*, conservation, microsatellite markers

*Ceratopteris thalictroides* (L.) Brongn. (Parkeriaceae) is a semi-aquatic fern with circum-tropical distribution, mainly growing in paddy fields, ponds, or marshes (Watano and Masuyama, 1994). This genus contains four species (Lloyd, 1974). Whereas three of those species are diploid plants, *C. thalictroides* is tetraploid (Hickok, 1977; Masuyama and Watano, 2005). Although *C. thalictroides* is widely distributed throughout tropical and subtropical regions, it is internationally considered to be of at least some concern, based on the IUCN Red List of Threatened Species.

In Korea, this species was first recorded at Suncheon and Gwangyang, in the southern regions of the Peninsula. Since then, more populations have been identified. This species is now designated as endangered in Korea and has been assigned legal protection for eight extant populations (Seocheon, Gunsan, Jeongeup, Buan, Iksan, Gochang, Gwangju, and Busan). These Korean populations are clearly critical because they represent the northern edge of the distribution range, possibly providing potential for further expansion of the species if those plants are able to adapt to selection pressures from such marginal environments (Kawecki, 2008). Therefore,

efforts to conserve Korean populations are needed if we are to achieve long-term survival for *C. thalictroides*.

In China, researchers are also taking various steps to conserve the populations of *C. thalictroides*. There, this species has been designated as endangered in an effort to protect and manage its native habitats (Yu, 1999). In addition, its genetic variability has been analyzed by using various markers such as random amplified polymorphic DNA, inter-simple sequence repeat, and microsatellites (Dong et al., 2008; Yang et al., 2016). Although 30 microsatellite markers have been established for Chinese populations of *C. thalictroides* (Yang et al., 2016), they are not adequate for frequently examining genetic polymorphism. Furthermore, it is cost-effective for such markers to be organized as “sets” if researchers are to continue periodic genetic monitoring. Therefore, we have developed new polymorphic microsatellite sets for evaluating the genetic status of Korean populations of *C. thalictroides* and conserving those plants.

### Materials and Methods

For these new microsatellite markers, total genomic DNA

\*Author for correspondence: quercus@jnu.ac.kr

was obtained from a fresh leaf collected from an individual plant of *C. thalictroides* growing at the Dongnim reservoir, Gochang, Korea. The extraction procedure involved an MG Plant Genomic DNA Extraction SV Miniprep Kit used according to the manufacturer's protocol (Macrogen Inc., Seongnam, Korea). After the quality of the genomic DNA was checked by gel electrophoresis, we generated a shotgun library by using the Illumina Mi-Seq platform (LAS, Seoul, Korea). In all, 5,256,220 paired-ends with read lengths of approximately 300 bp were obtained. Based on screening with the SSR\_Pipeline v.0951 infrastructure (Miller et al., 2013), we identified a 22,990 microsatellite motif with flanking regions larger than 100 bp that contained di-, tri-, or tetra-nucleotide repeats that were at least 12, 6, or 5 times, respectively, in size. Following the method of Cho et al. (2015), we assembled groupings of 50 to 100 reads that were reference guide-mapped on the whole raw reads, using the Geneious R10.1.3 software platform (Kearse et al., 2012). From this, we selected reads that had two separate alleles and no additional single nucleotide polymorphisms in the flanking region. After eliminating any duplication through *de novo* assembly, we used the final chosen reads to develop microsatellite markers. We then utilized the Primer 3 software in Geneious program R10.1.3 to design 81 primer pairs (18–22 bp long), using a melting temperature of 53–60°C and GC contents of 35–65%. The forward primers added three sets of M13 tag sequences (5'-CACGACGTTGTAAACGAC-3', 5'-TGTGGAATTGTGAGCGG-3', and 5'-CTATAGGGCACGCGTGGT-3') with 6-FAM, VIC, and NED fluorescent dye, respectively. For multiplex polymerase chain reaction (PCR), each primer set comprised nine primers with lengths of 100–300 bp.

To test the effectiveness of these new microsatellite loci, we sampled 63 individuals of *C. thalictroides* from two

populations (Dongnim reservoir, Gochang,  $n = 33$ ; Is. Doonchido, Busan,  $n = 30$ ). For each locus, PCR amplification was performed in a final volume of 5  $\mu$ L that contained 15 to 20 ng of extracted DNA, 2.5  $\mu$ L of 2 $\times$  Multiplex PCR master mix (Qiagen, Valencia, CA, USA), 0.01  $\mu$ M for the forward primer, 0.2  $\mu$ M for the reverse primer, and 0.1  $\mu$ M for the M13 primer. The PCR protocol included an initial denaturation for 15 min at 95°C; followed by 30 cycles, each consisting of denaturing for 30 s at 94°C, 1.5 min of annealing at 52°C, and extension for 1.5 min at 72°C; with a final extension for 10 min at 72°C. The PCR products were analyzed using an ABI 3730XL sequencer with GeneScan-500LIZ Size Standard (Applied Biosystems, Foster City, CA, USA). Allele sizes and the peaks of each sample were determined via Peak Scanner software 2 (Applied Biosystems). In analyzing the tetraploid microsatellite data, we were not able to make exact determinations of allele frequencies and patterns due to the uncertainty of multiple alleles (Dufresne et al., 2014). Therefore, we calculated the number of alleles ( $N_A$ ), the Shannon-Wiener Diversity Index ( $H'$ ), and the expected heterozygosity ( $H_E$ ) using Atetra 1.2 software (van Puyvelde et al., 2010), which was developed for analyzing codominant microsatellite data in tetraploid species.

## Results and Discussion

We have produced useful sets of microsatellite markers for analyzing genetic diversity and establishing conservation strategies for *Ceratopteris thalictroides*, an endangered plant. Of the 81 primer pairs studied here, 25 microsatellite markers were polymorphic, and 17 were found to be monomorphic in two of the examined populations (Table 1). All developed markers have been deposited in GenBank (Table 1). Among the 25 polymorphic microsatellite loci, the number of alleles per locus

**Table 1.** Characterization of 25 microsatellite loci for *Ceratopteris thalictroides*.

Locus <sup>a</sup>	Primer sequence (5'–3')	Repeat motif	$N_A$	Allele size range (bp)	Fluorescent label	GenBank accession No.
		Multiplex mix A				
cet007	F: AAAATACTGGCCACGGTTG R: AAGATGTTGAAGTGGGCTG	(AG) <sub>17</sub>	8	153–179	NED	MH261329
cet009	F: TGAATTGCTCATGATGTTGC R: AGACTCGTTTTTCATGGAGAC	(GAA) <sub>7</sub>	3	232–241	NED	MH261330
cet017	F: CAAGAGGAGGCAGAATAACA R: TCTCTCTCCTAGGTTTCAAGT	(AAG) <sub>16</sub>	5	210–222	NED	MH261333
cet023	F: CCTCACTCTCTTTCCAAACT R: CATTCCGATACACCAAGCTA	(CTC) <sub>7</sub>	5	180–207	VIC	MH261337
cet028*	F: ACATGCTTCTTTGATCTCGT R: TATGATCTTTGATGCAGGCA	(AG) <sub>12</sub>	4	179–189	6-FAM	MH261339

Table 1. Continued.

Locus <sup>a</sup>	Primer sequence (5'–3')	Repeat motif	$N_A$	Allele size range (bp)	Fluorescent label	GenBank accession No.
cet101	F: ACTTTGAACTATTCTGCATCG R: TTGTTGTACATTTGTTGGCT	(AC) <sub>13</sub>	2	98–102	6-FAM	MH261347
cet104	F: TGCCATAACAAATGTCAAGA R: CATTACACTATGCACATGGA	(TACA) <sub>5</sub>	2	139–151	VIC	MH261348
cet120	F: GTCAGAATCCCGTCAGTATC R: CAACAAGAGGGACAGCTTAT	(GTAT) <sub>9</sub>	5	222–238	6-FAM	MH261352
Multiplex mix B						
cet010*	F: TAAAAGCGATCTCCACGTAG R: TAAAGCTTGGAGAGAGGCTA	(AG) <sub>16</sub>	9	143–165	6-FAM	MH261331
cet012*	F: AGTCAAAACTCTGTGCTAGG R: TACGTTGGCTATGTTCCCTC	(AG) <sub>13</sub>	5	227–241	6-FAM	MH261332
cet018*	F: CAGGATAGAGAGGATTCGC R: AGGAGGAGCATTCTATGACT	(TC) <sub>24</sub>	7	242–254	NED	MH261334
cet022	F: TCATATCGCAATCAAAGCAG R: TGGAAGAGAGAGCAATTG	(TC) <sub>17</sub>	7	177–189	VIC	MH261336
cet025	F: GGAGAGAAATGCTTGTATTGT R: ATGACGATCATGAATGGGTG	(TG) <sub>12</sub>	8	175–189	NED	MH261338
cet042*	F: GAAGACATCACCTCCTCTG R: CAAGAAAGAGCAAGGAGTCT	(GA) <sub>13</sub>	3	267–271	VIC	MH261342
cet046	F: GACACCATTCATGCGAC R: ATACTTGCCTGTGTGAGAG	(CT) <sub>12</sub>	3	185–189	6-FAM	MH261343
cet050	F: CAATGAGCAGAGTTGTGAAG R: TCATGGTTGTTTTGGAGGAA	(CT) <sub>17</sub>	9	209–233	VIC	MH261344
cet108	F: CCATCAATTTGAGTCGAGGAT R: ACAGAGTTGCACAAGGTATT	(TG) <sub>12</sub>	5	161–171	NED	MH261350
Multiplex mix C						
cet021*	F: GCAACAATAAAGCGTCAAT R: TGGTCACTGACGAATCAAAT	(AC) <sub>16</sub>	6	240–254	6-FAM	MH261335
cet029*	F: TACAGTGACAATGCTTTCCT R: CTCTGAGCCTTCCTTTTCTT	(GAA) <sub>6</sub>	4	213–222	6-FAM	MH261340
cet032*	F: TTCTTCTCAGGCACCTTTT R: AGTTCTGAGAGTCCACAATG	(CAA) <sub>6</sub>	3	209–215	VIC	MH261341
cet051	F: GACGATGGAGGCATTATGAT R: GATAGGTTCTAGGCGCATT	(GA) <sub>22</sub>	14	291–315	VIC	MH261345
cet053	F: CTTGGGATGCGAGAAATAGT R: CCATCCTCATCTTCACCAAA	(TC) <sub>15</sub>	7	226–242	NED	MH261346
cet107	F: GGCCTCTGTGGAATATGATT R: TTCATGTCCTAACTCAACCG	(TG) <sub>17</sub>	3	158–162	NED	MH261349
cet113	F: GTCCCGACTTAAATCCCAT R: GCAAAATTGTTGGGCAGAC	(CA) <sub>15</sub>	4	169–187	VIC	MH261351
cet127	F: TTGAAGTTGGAGCATGAAAG R: TGGCGAATATGACATACCTT	(CT) <sub>16</sub>	9	257–275	NED	MH261353

$N_A$ , number of alleles.

<sup>a</sup>Reaction concentrations in PCR for primers were 0.01 for forward primer and 0.2 for reverse primer. Loci marked with asterisk (\*) had reaction concentrations of 0.02 for forward primer and 0.4 for reverse primer.

**Table 2.** Summary of genetic parameters estimated from 25 microsatellite loci across 63 individuals sampled from the two Korean populations, Gochang (GC) and Busan (BS), Korea.

Locus	GC ( $n = 33$ )				BS ( $n = 30$ )			
	$N_A$	$H_E$	$H'$	$Sr$	$N_A$	$H_E$	$H'$	$Sr$
cet007	2	0.114	0.229	153–155	8	0.826	1.856	153–179
cet009	3	0.621	1.021	232–241	3	0.583	0.946	232–241
cet010	7	0.718	1.425	143–163	8	0.810	1.825	145–165
cet012	4	0.501	0.861	227–237	4	0.589	1.084	227–241
cet017	5	0.685	1.307	210–222	2	0.320	0.500	213–222
cet018	6	0.670	1.322	242–254	5	0.627	1.223	244–254
cet021	5	0.622	1.190	240–254	4	0.643	1.130	240–252
cet022	6	0.637	1.297	179–189	4	0.266	0.548	177–187
cet023	4	0.691	1.244	180–207	4	0.692	1.256	180–207
cet025	8	0.784	1.733	175–189	7	0.749	1.541	177–189
cet028	3	0.601	0.994	179–189	3	0.604	1.010	179–189
cet029	4	0.729	1.339	213–222	4	0.728	1.341	213–222
cet032	2	0.497	0.691	209–215	3	0.564	0.906	209–215
cet042	2	0.114	0.229	267–269	3	0.499	0.693	267–271
cet046	2	0.165	0.305	187–189	3	0.615	1.025	185–189
cet050	9	0.717	1.495	209–233	6	0.785	1.620	209–231
cet051	11	0.859	2.124	295–315	13	0.860	2.152	291–313
cet053	7	0.687	1.414	226–242	6	0.643	1.324	226–240
cet101	2	0.030	0.077	98–102	1	0.000	0.000	98
cet104	2	0.497	0.690	139–151	2	0.453	0.645	139–151
cet107	2	0.030	0.077	160–162	3	0.183	0.368	158–162
cet108	5	0.730	1.424	161–171	4	0.679	1.203	161–169
cet113	2	0.497	0.691	169–185	4	0.637	1.163	169–187
cet120	4	0.680	1.224	226–238	3	0.452	0.731	222–234
cet127	5	0.640	1.208	265–273	7	0.803	1.771	257–275

$N_A$ , number of alleles;  $H_E$ , expected heterozygosity;  $H'$ , Shannon-Wiener Diversity Index;  $Sr$ , size range (bp).

ranged from 2 to 13, with an average of 5.64. Values for ( $H_E$ ) were 0.000 to 0.860, while  $H'$  ranged from 0.000 to 2.152 (Table 2). These results are similar to those reported from a previous study in China (Yang et al., 2016) that involved RAD tag sequencing ( $N_A$ , 4–10;  $H_E$ , 0.264–0.852). However, because the Korean Peninsula is a northern limit for the distribution of *C. thalictroides*, our findings suggest that the marker sets described here include considerable polymorphism.

### Acknowledgments

This work was supported by the National Institute of Biological Resources of Korea (Grant No. NIBR201703101).

### Conflict of Interest

Authors declare that there is no conflict of interest.

### Literature Cited

- Cho, W.-B., I.-S. Choi and B.-H. Choi. 2015. Development of microsatellite markers for the endangered *Pedicularis ishidoyana* (Orobanchaceae) using next-generation sequencing. *Applications in Plant Sciences* 3: 1500083.
- Dong, Y.-H., J.-M. Chen, G. W. Robert and Q.-F. Wang. 2008. Genetic variation in the endangered aquatic fern *Ceratopteris thalictroides* (Parkeriaceae) in China: implications from

- RAPD and ISSR data. *Botanical Journal of the Linnean Society* 157: 657–671.
- Dufresne, F., M. Stift, R. Vergilino and B. K. Mable. 2014. Recent progress and challenges in population genetics of polyploid organisms: an overview of current state-of-the-art molecular and statistical tools. *Molecular Ecology* 23: 40–69.
- Hickok, L. G. 1977. Cytological relationships between three diploid species of the ferns genus *Ceratopteris*. *Canadian Journal of Botany* 55: 1660–1667.
- Kawecki, T. J. 2008. Adaptation to marginal habitats. *Annual Review of Ecology, Evolution, and Systematics* 39: 321–342.
- Kearse, M., R. Moir, A. Wilson, S. Stones-Havas, M. Cheung, S. Sturrock, S. Buxton, A. Cooper, S. Markowitz, C. Duran, T. Thierer, B. Ashton, P. Meintjes and A. Drummond. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28: 1647–1649.
- Lloyd, R. M. 1974. Systematic of the genus *Ceratopteris* Brongn. (Parkeriaceae) II. Taxonomy. *Brittonia* 26: 139–160.
- Masuyama, S. and Y. Watano. 2005. Cryptic species in the fern *Ceratopteris thalictroides* (L.) Brongn. (Parkeriaceae). II. Cytological characteristics of three cryptic species. *Acta Phytotaxonomica et Geobotanica* 56: 231–240.
- Miller, M. P., B. J. Knaus, T. D. Mullins and S. M. Haig. 2013. SSR\_pipeline: a bioinformatic infrastructure for identifying microsatellites from paired-end Illumina high-throughput DNA sequencing data. *Journal of Heredity* 104: 881–885.
- van Puyvelde, K., A. van Geert and L. Triest. 2010. ATETRA, a new software program to analyse tetraploid microsatellite data: comparison with TETRA and TETRASAT. *Molecular Ecology Resources* 10: 331–334.
- Watano, Y. and S. Masuyama. 1994. Genetic differentiation in populations of the polymorphic fern *Ceratopteris thalictroides* in Japan. *Journal of Plant Research* 107: 139–146.
- Yang, X. Y., Z. C. Long, A. W. Gichira, Y. H. Guo, Q. F. Wang and J. M. Chen. 2016. Development of microsatellite markers in the tetraploid fern *Ceratopteris thalictroides* (Parkeriaceae) using RAD tag sequencing. *Genetics and Molecular Research* 15: gmr7550.
- Yu, Y. F. 1999. A milestone of wild plant conservation in China. *Plants* 5: 3–11.